

CORRECTION

Open Access



Correction: Tubulointerstitial nephropathy is the predominant finding in men in a review of more than 3000 renal biopsies over a 10-year period from Sri Lanka

Jennifer Pett^{1,2}, Christine Linhart^{1*}, Nicholas Osborne^{1,3*}, Stephen Morrell¹, Mohammed Fahim^{4,5}, John Knight^{1,2}, Shakila Premarathne^{6,7}, A. W. M. Wazil⁸, Neelakanthi Ratnatunga⁹, Sulcochana Wijethunga⁹, Shenal Thalgahagoda¹⁰, Zoltan Endre^{5,11}, Richard Taylor¹ and Nishantha Nanayakkara⁸

Correction: BMC Nephrology (2024) 25:295
<https://doi.org/10.1186/s12882-024-03731-7>

Following publication of the original article [1], the authors identified that the references within the article were not listed sequentially.

The online version of the original article can be found at <https://doi.org/10.1186/s12882-024-03731-7>.

*Correspondence:

Christine Linhart
c.linhart@unsw.edu.au

Nicholas Osborne
n.osborne@uq.edu.au

¹School of Population Health, University of New South Wales, UNSW, Sydney, Australia

²The George Institute for Global Health, Sydney, Australia

³School of Public Health, University of Queensland, Herston, Australia

⁴South Asian Clinical Toxicology Research Collaboration, Faculty of Medicine, University of Peradeniya, Peradeniya, Sri Lanka

⁵School of Clinical Medicine, Faculty of Medicine and Health, UNSW, Sydney, Australia

⁶Department of Microbiology, Faculty of Medicine, University of Peradeniya, Peradeniya, Sri Lanka

⁷Centre for Research, Nephrology and Kidney Transplant Unit, National Hospital, Kandy, Sri Lanka

⁸Nephrology and Kidney Transplant Unit, National Hospital, Kandy, Sri Lanka

⁹Department of Pathology, Faculty of Medicine, University of Peradeniya, Peradeniya, Sri Lanka

¹⁰Department of Paediatrics, Faculty of Medicine, University of Peradeniya Teaching Hospital, Peradeniya, Sri Lanka

¹¹Department of Nephrology, Prince of Wales Hospital, Randwick, Sydney, Australia

Also, they would like to correct the below two errors:

1. Nugegoda was listed as west of Colombo, this has been changed to east.
2. The proportion of TIN in the study by Pilapitiya et al. was incorrectly listed as 25.5, this has been changed to 27.1.

The original article has been corrected.

Published online: 28 October 2024

References

1. Pett et al. BMC Nephrol. (2024);25:295 <https://doi.org/10.1186/s12882-024-03731-7>

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.